



ELSEVIER

Journal of Pure and Applied Algebra 146 (2000) 35–44

JOURNAL OF
PURE AND
APPLIED ALGEBRA

www.elsevier.com/locate/jpaa

on and similar papers at core.ac.uk

provi

On weak π -regularity of rings whose prime ideals are maximal

Chan Yong Hong^{a,*}, Nam Kyun Kim^a, Tai Keun Kwak^b, Yang Lee^c

^a*Department of Mathematics, Kyung Hee University, Seoul 130-701, South Korea*

^b*Department of Mathematics, Daejin University, Pocheon 487-800, South Korea*

^c*Department of Mathematics, Busan National University, Busan 609-735, South Korea*

Communicated by G.M. Kelly; received 29 September 1997

Abstract

We investigate, in this paper, the connections between the weak π -regularity and the maximality of prime ideals in 2-primal rings, right quasi-duo rings and PI-rings, respectively. © 2000 Elsevier Science B.V. All rights reserved.

MSC: 16E50; 16D30; 16U80

The relationship between various generalizations of von Neumann regularity and the condition that every prime ideal is maximal have been investigated by many authors [3, 4, 7, 9, 13, 15, 16]. The first clearly established equivalence between a generalization of von Neumann regularity and the maximality of prime ideals seems to have been made by Storrer [13] in the following result: If R is a commutative ring with identity then R is π -regular if and only if every prime ideal of R is maximal. Storrer's result was extended to PI-rings [7, Theorem 2.3], right duo rings [9, Corollary 1] and bounded weakly right duo rings [15, Theorem 3], respectively. On the other hand, Hirano [9] proved that if R is a 2-primal ring, then R is π -regular if and only if every prime ideal of R is a maximal one-sided ideal. Recently, Birkenmeier et al. [3] showed that if R is a 2-primal ring, then $R/P(R)$ is right weakly π -regular if and only if every prime ideal of R is maximal. These results were mainly explained the relation between the π -regularity and the maximality of prime ideals of rings.

The π -regularity of rings is extended to the weak π -regularity. In general, π -regular rings are weakly π -regular rings but the converse is not hold.

* Corresponding author.

E-mail address: hcy@nms.kyunghee.ac.kr (C.Y. Yong)

We investigate the connections between the results of previously mentioned papers and weak π -regularity in 2-primal rings, right quasi-duo rings and PI-rings, respectively. Consequently, our results in this paper extend many of results in [4, 7, 15].

Throughout this paper the letter R denotes an associative ring with identity and all prime ideals of R are assumed to be proper. $P(R)$, $J(R)$ and $N(R)$ denote the prime radical, the Jacobson radical and the set of all nilpotent elements of R , respectively.

We begin with the following definitions.

Definition 1. (1) A ring R is said to be (*strongly*) π -regular if for every $x \in R$ there exists a natural number n , depending on x , such that $(x^n \in x^{n+1}R)x^n \in x^nRx^n$. Strong π -regularity is right–left symmetric [6].

(2) A ring R is said to be *right (left) weakly π -regular* if for every $x \in R$ there exists a natural number n , depending on x , such that $x^n \in x^nRx^nR$ ($x^n \in Rx^nRx^n$). R is *weakly π -regular* if it is both right and left weakly π -regular [8].

Definition 2. A ring R is called *2-primal* if $P(R) = N(R)$ [2].

The term *2-primal* was come upon originally by Birkenmeier, Heatherly and Lee. But Hirano [9] used the term *N -ring* for what we call a 2-primal ring. The 2-primal condition was taken up independently by Sun [14], where in the setting of rings he introduced a condition to be called *weakly symmetric*, which is equivalent to the 2-primal condition for rings.

Hirano showed the following proposition.

Proposition 3 (Hirano [9, Theorem 1]). *Let R be a 2-primal ring. Then the following statements are equivalent:*

- (a) R is strongly π -regular.
- (b) R is π -regular.
- (c) $R/J(R)$ is π -regular and $J(R)$ is nil.
- (d) Every prime ideal of R is a maximal one-sided ideal.

The following example shows that in a 2-primal ring “weak π -regularity” is not the same “ π -regularity”. Hence in Proposition 3, condition “(b)” cannot be replaced by the condition “ R is right weakly π -regular”.

Example 4. Let D be a simple domain which is not a division ring. We consider the ring

$$R = \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \mid a, b \in D \right\}.$$

Then

(1) it can be easily checked that R is 2-primal with

$$P(R) = \begin{pmatrix} 0 & D \\ 0 & 0 \end{pmatrix}.$$

(2) R is right weakly π -regular: If $x \in P(R)$, then $x^2 = 0$ and hence $x^2 \in x^2 Rx^2 R$. Suppose $x \notin P(R)$. Then since $R/P(R) \cong D$ is simple, $RxR + P(R) = R$. Thus $\alpha + \beta = 1$ for some $\alpha \in RxR$ and $\beta \in P(R)$. But $\beta^2 = 0$ and so $1 = (\alpha + \beta)^2 \in RxR$. Hence $x \in xRxR$. Therefore R is right weakly π -regular.

(3) R is not π -regular: Let $a \neq 0$ and 1. Then

$$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}^n \notin \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}^n R \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}^n$$

for any positive integer n .

As a parallel result to this Proposition 3, we obtain the following result.

Proposition 5. *Let R be a 2-primal ring. Then the following statements are equivalent:*

- (a) $R/J(R)$ is right weakly π -regular and $J(R)$ is nil.
- (b) Every prime ideal of R is maximal.

Proof. (a) \Rightarrow (b): Assume (a). Since $J(R)$ is nil and R is 2-primal, $J(R) = P(R)$. By [3, Corollary 9], every prime ideal of R is maximal.

(b) \Rightarrow (a): Assume (b). By [3, Corollary 9], $R/P(R)$ is right weakly π -regular and so $R/J(R)$ is right weakly π -regular. Now let $a \in J(R)$. Consider $\bar{a} \in \bar{R} = R/P(R)$. Since \bar{R} is right weakly π -regular, there exists a positive integer n such that $\bar{a}^n = \bar{a}^n \bar{r}$ for some $\bar{r} \in \bar{R} \bar{a}^n \bar{R} \subseteq \bar{J}(R)$, where $\bar{J}(R) = J(R)/P(R)$. Then $\bar{a}^n(1 - \bar{r}) = \bar{0}$ and so $a^n \in P(R)$. Since R is 2-primal, $a \in P(R)$ and hence $J(R)$ is nil. \square

Related to this Proposition 5, we noted that there exists a 2-primal ring whose prime ideals are maximal but neither right nor left weakly π -regular [3, Example 12].

However, we have the following theorem.

Theorem 6. *Let R be a 2-primal ring whose primitive factor rings are Artinian. Then the following statements are equivalent:*

- (a) R is strongly π -regular.
- (b) R is π -regular.
- (c) R is weakly π -regular.
- (d) R is right weakly π -regular.
- (e) $R/J(R)$ is right weakly π -regular and $J(R)$ is nil.
- (f) Every prime ideal of R is maximal.

Proof. It suffices to show that (f) \Rightarrow (a). Assume (f). Let P be a prime ideal of R . Then P is maximal, so it is primitive. Thus R/P is Artinian. Hence R/P is strongly π -regular and so R is strongly π -regular by [7, Theorem 2.1]. \square

Recall that a ring R is said to be *right (left) quasi-duo* if every maximal right (left) ideal of R is two sided. A ring R is said to be *of bounded index (of nilpotency)* if there exists a positive integer n such that $a^n = 0$ for all nilpotent elements a of R .

The following theorem extends [4, Proposition 2.14].

Theorem 7. *Let R be a right quasi-duo ring. Then the following statements are equivalent:*

- (a) R is right weakly π -regular.
- (b) R is strongly π -regular.

Proof. It is enough to show that (a) \Rightarrow (b). Assume (a). Then for any $a \in R$ there exists a positive integer n such that $a^n R = a^n R a^n R$. We claim that $aR + r(a^n) = R$, where $r(a^n)$ is the right annihilator of a^n . If not, there is a maximal right ideal M of R such that $aR + r(a^n) \subseteq M$. Then $a^n R = a^n M$, and so $a^n = a^n b$ for some $b \in M$. Hence $a^n(1 - b) = 0$ and so $1 - b \in r(a^n) \subseteq M$, which is a contradiction. Therefore R is strongly π -regular. \square

The fact that a right duo ring (i.e., a ring whose right ideals are two sided) implies 2-primal lead one to conjecture that a right quasi-duo ring is 2-primal. However in the following examples, the rings R are right quasi-duo but not 2-primal. Moreover, these examples show that the conditions of the following Proposition 12 and Corollary 13 are required.

Example 8. Let T be the $n \times n$ upper triangular matrix ring over a field F , where n is an infinite cardinal number. Then by [16, Proposition 2.1] T is right quasi-duo. Now consider $R = T[[x]]$, where $T[[x]]$ denotes the ring of formal power series over T . Let M' be a maximal right ideal of R . Note that M' is generated by x and a maximal right ideal M of T . Since T is right quasi-duo, M' is a two-sided ideal of R . Thus R is also right quasi-duo.

Next, we claim that R is not 2-primal. Let e_{ij} be the infinite matrix over F with (i, j) -entry the identity 1 and elsewhere the zero element 0. Now in the formal power series ring R , let

$$f(x) = e_{12}x + (e_{34} + e_{56})x^2 + \cdots + \left(\sum_{i=0}^{2^{n-1}-1} e_{(2^n+2i-1)(2^n+2i)} \right) x^n + \cdots$$

and

$$g(x) = e_{23}x + (e_{45} + e_{67})x^2 + \cdots + \left(\sum_{i=0}^{2^{n-1}-1} e_{(2^n+2i)(2^n+2i+1)} \right) x^n + \cdots .$$

Then $f(x)^2 = 0$ and $g(x)^2 = 0$. But

$$f(x) + g(x) = (e_{12} + e_{23})x + (e_{34} + e_{45} + e_{56} + e_{67})x^2 + \cdots \\ + \left(\sum_{i=0}^{2^n-1} (e_{(2^n+2i-1)(2^n+2i)} + e_{(2^n+2i)(2^n+2i+1)}) \right) x^n + \cdots$$

and by setting

$$\alpha_n = \sum_{i=0}^{2^n-1} (e_{(2^n+2i-1)(2^n+2i)} + e_{(2^n+2i)(2^n+2i+1)}),$$

we have $\alpha_n^{2^n-1} \neq 0$ and $\alpha_n^{2^n} = 0$. Thus $f(x) + g(x)$ is not nilpotent; whence $f(x) \notin P(R)$ or $g(x) \notin P(R)$. Therefore R is not 2-primal. Moreover R is of no bounded index and $J(R)$ is not nil.

Example 9. Let F be a field and S denote the full ring of 2×2 matrices over F and

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in S \mid a \in F \right\}.$$

We consider the ring $R = T + xS[[x]]$, where $S[[x]]$ denotes the ring of formal power series over S . Then $xS[[x]]$ is the unique maximal left and right ideal of R . Thus R is a right quasi-duo ring and $J(R) = xS[[x]]$ is not nil. Moreover, R is a semiprime PI-ring of bounded index 2. But R is not 2-primal, since

$$x \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

is a nonzero nilpotent element.

Example 10 (Birkenmeier et al. [4, Example 3.3]). Let G be an abelian group which is the direct sum of a countably infinite number of infinite cyclic groups; and denote by $\{b(0), b(1), b(-1), \dots, b(i), b(-i), \dots\}$ of basis of G . Then there exists one and only one homomorphism $u(i)$ of G , for $i = 1, 2, \dots$ such that $u(i)(b(j)) = 0$ if $j \equiv 0 \pmod{2^i}$ and $u(i)(b(j)) = b(j-1)$ if $j \not\equiv 0 \pmod{2^i}$. Denote U the ring of endomorphisms of G generated by the endomorphisms $u(1), u(2), \dots$. Now, let A be the ring obtained from U by adjoining the identity map of G . Let $R = A \otimes_{\mathbb{Z}} \mathbb{Q}$, where \mathbb{Z} is the ring of integers and \mathbb{Q} is the field of rationals. Then R is a right quasi-duo ring of no bounded index and $J(R)$ is nil but R is not 2-primal. R is also a semiprime strongly π -regular ring, but there exists a prime ideal of R which is not maximal. If every prime ideal of R is maximal, then $J(R) = P(R) = 0$, which is a contradiction.

Now, we show that a right quasi-duo ring R of bounded index with $J(R)$ nil is 2-primal.

Lemma 11. *Let I be a right ideal of R and n a positive integer. If $a^n = 0$ for all $a \in I$, then $a^{n-1}Ra^{n-1} = 0$.*

Proof. For any $r \in R$, let $b = a^{n-1}r \in I$. Then $ab = a(a^{n-1}r) = a^n r = 0$. Thus $(b+a)^n = b^n + b^{n-1}a + b^{n-2}a^2 + \cdots + ba^{n-1} + a^n$. Since $(b+a)^n = 0$ and $a^n = 0 = b^n$, then we have

$$\begin{aligned} 0 &= b^{n-1}a + b^{n-2}a^2 + \cdots + b^2a^{n-2} + ba^{n-1} \\ &= b(a^{n-1}r)b^{n-3}a + b(a^{n-1}r)b^{n-4}a^2 + \cdots + b(a^{n-1}r)a^{n-2} + ba^{n-1} \\ &= ba^{n-1}(sa + 1) \quad \text{for some } s \in R. \end{aligned}$$

Since $(sa)^{n+1} = s(as)^na = 0$, $sa + 1$ is invertible. Thus $ba^{n-1} = 0$ and hence $a^{n-1}ra^{n-1} = 0$. Therefore $a^{n-1}Ra^{n-1} = 0$. \square

Proposition 12. *Let R be a right quasi-duo ring of bounded index with $J(R)$ nil. Then R is a 2-primal ring.*

Proof. Since R is right quasi-duo and $J(R)$ is nil, $N(R) = J(R)$ by [16, Lemma 2.3]. Let m be the bounded index of R . If $m = 1$, then $P(R) = N(R) = 0$. Let $m \geq 2$ and $J(R) \neq 0$. Then $J(R) = P(R)$. If not, then $J(R)/P(R)$ is a nonzero nil ideal of $\bar{R} = R/P(R)$ with bounded index $n \geq 2$. Thus there exists a nonzero $\bar{a} \in J(R)/P(R)$ such that $\bar{a}^{n-1} \neq \bar{0}$. By Lemma 11, $\bar{a}^{n-1}\bar{R}\bar{a}^{n-1} = \bar{0}$ and so $(\bar{R}\bar{a}^{n-1}\bar{R})^2 = \bar{0}$. Since \bar{R} is semiprime, $\bar{R}\bar{a}^{n-1}\bar{R} = \bar{0}$, which is a contradiction. Therefore R is 2-primal. \square

Note that in Proposition 12 the conditions “ R is of bounded index” and “ $J(R)$ is nil” are not superfluous (Examples 9 and 10).

Corollary 13. *Let R be a right quasi-duo ring of bounded index. Then the following statements are equivalent:*

- (a) R is strongly π -regular.
- (b) R is π -regular.
- (c) $R/J(R)$ is π -regular and $J(R)$ is nil.
- (d) R is weakly π -regular.
- (e) R is right weakly π -regular.
- (f) $R/J(R)$ is right weakly π -regular and $J(R)$ is nil.
- (g) $R/P(R)$ is right weakly π -regular.
- (h) Every prime ideal of R is maximal.

Proof. (a) \Leftrightarrow (b) \Leftrightarrow (d) \Leftrightarrow (e) and (c) \Leftrightarrow (f): These follow from Theorem 7.

(e) \Rightarrow (f) and (e) \Rightarrow (g): These are obvious.

(g) \Rightarrow (a): Assume (g). Then $R/P(R)$ is strongly π -regular by Theorem 7. Thus R is strongly π -regular by [7, Theorem 2.1].

(f) \Rightarrow (h): Assume (f). Then by Proposition 12 and Proposition 5, every prime ideal of R is maximal.

(h) \Rightarrow (a): It follows from [16, Theorem 2.5]. \square

In Corollary 13, the condition “ R is of bounded index” is not superfluous (Example 10).

Remark. In Example 4, the ring R is 2-primal right weakly π -regular which is of bounded index 2. But it is not π -regular. The ring in [3, Example 12] is a 2-primal ring whose prime ideals are maximal, and it is of bounded index 2. But it is neither right nor left weakly π -regular.

Recall that a ring R is said to be *weakly right duo* if for every $a \in R$ there is a positive integer $n = n(a)$, depending on a , implies $a^n R$ is a two-sided ideal of R [15]. A ring R is said to be *abelian* if every idempotent element of R is central. Note that every weakly right duo ring is abelian right quasi-duo [16, Proposition 2.2] but, in general, is of no bounded index. For example, the ring R in Example 10 is weakly right duo but it is of no bounded index.

However, we have the following proposition.

Proposition 14. *Let R be a weakly right duo ring. Then the following statements are equivalent:*

- (a) R is strongly π -regular.
- (b) R is π -regular.
- (c) R is right (left) weakly π -regular.
- (d) $R/J(R)$ is strongly regular.

Proof. It is sufficient to show that (c) \Rightarrow (d) and (d) \Rightarrow (a).

(c) \Rightarrow (d): Assume (c). Then $R/J(R)$ is strongly π -regular by Theorem 7. Also by [15, Corollary 2], $R/J(R)$ is reduced and so it is strongly regular.

(d) \Rightarrow (a): It follows from [15, Corollary 2; 1, Theorem 3] and Theorem 7. \square

In weakly right duo rings, if every prime ideal of R is maximal then R is strongly π -regular [16, Theorem 2.5]. However, the converse is not true (Example 10).

Now we investigate the connection between the weak π -regularity and the maximality of prime ideals in PI-rings.

Theorem 15. *Let R be a prime right Goldie ring. If R is right weakly π -regular, then R is a simple ring.*

Proof. Let I be a nonzero two-sided ideal of R . Since R is prime, I is essential as a right ideal. Then by [5, Theorem 1.10], I contains a regular element $a \in R$. Since R is

right weakly π -regular, there exists a positive integer n such that $a^n R = a^n R a^n R$ and so $a^n = a^n b$ for some $b \in R a^n R$. Thus $a^n(1 - b) = 0$. Now a^n is also regular element and so $b = 1$; whence $I = R$. Therefore R is a simple ring. \square

Corollary 16. *Let R be a PI-ring. If R is right weakly π -regular, then every prime ideal of R is maximal.*

Proof. Let P be a prime ideal of R . Then R/P is a prime PI-ring. So by Posner's Theorem [12], R/P is right Goldie. By Theorem 15, R/P is simple. Hence every prime ideal of R is maximal. \square

Theorem 17. *Let R be a PI-ring. Then the following statements are equivalent:*

- (a) R is right (left) weakly π -regular.
- (b) $R/J(R)$ is right (left) weakly π -regular and $J(R)$ is nil.

Proof. (b) \Rightarrow (a): Assume (b). If R is not right weakly π -regular, then R is not strongly π -regular. Thus there exists a prime ideal P of R such that R/P is not strongly π -regular by [7, Theorem 2.1]. Since $R/J(R)$ is a right weakly π -regular and PI-ring, $R/J(R)$ is strongly π -regular by [9, Theorem 4]. If $J(R) \subseteq P$, then a homomorphic image R/P of $R/J(R)$ is strongly π -regular, which is a contradiction. Thus $J(R) \not\subseteq P$ and so $(J(R) + P)/P$ is a nonzero two-sided ideal of R/P . Since R/P is a prime PI-ring, and so it is right Goldie, $(J(R) + P)/P$ contains a regular element $a + P$ with $a \in J(R)$ by [5, Theorem 1.10]. However, $J(R)$ is nil by assumption so $a^k = 0$ for some integer k ; whence $(a + P)^k = 0$ in R/P , which is also a contradiction. Therefore R is right weakly π -regular. \square

As a byproduct of Corollary 16 and Theorem 17, we obtain Corollary 18 which includes [7, Theorem 2.3; 9, Theorem 4].

Corollary 18. *Let R be a PI-ring. Then the following statements are equivalent:*

- (a) R is strongly π -regular.
- (b) R is π -regular.
- (c) $R/P(R)$ is π -regular.
- (d) $R/J(R)$ is π -regular and $J(R)$ is nil.
- (e) R is weakly π -regular.
- (f) R is right (left) weakly π -regular.
- (g) $R/J(R)$ is right weakly π -regular and $J(R)$ is nil.
- (h) Every prime factor ring of R is right (left) weakly π -regular.
- (i) Every prime ideal of R is maximal.
- (j) Every prime factor ring of R is simple Artinian.

Note that all primitive factor rings of a PI-ring and a right quasi-duo ring are Artinian.

Related to Corollaries 13 and 18, we have the following proposition.

Proposition 19. *Let R be of bounded index whose primitive factor rings are Artinian. Then the following statements are equivalent:*

- (a) R is strongly π -regular.
- (b) R is π -regular.
- (c) $R/P(R)$ is π -regular.
- (d) Every prime ideal of R is maximal.

Proof. (a) \Rightarrow (b) \Rightarrow (c): It is clear.

(c) \Rightarrow (d): Assume (c). Since R is of bounded index, $R/P(R)$ is also of bounded index. Then by [10, Proposition 2], every prime ideal of $R/P(R)$ is maximal. Therefore every prime ideal of R is maximal.

(d) \Rightarrow (a): Assume (d). Let P be a prime ideal of R . Then R/P is Artinian and so R/P is strongly π -regular. Thus by [7, Theorem 2.1], R is strongly π -regular. \square

The conditions of Proposition 19(1) “ R is of bounded index” and (2) “every primitive factor ring of R is Artinian” are not superfluous.

Example 20. (1) In Example 10, the ring R is a right quasi-duo ring, so every primitive factor ring of R is Artinian. R is also a strongly π -regular ring of no bounded index. But there exists a prime ideal of R which is not maximal.

(2) [3, Example 13]. Let $W = W_1[F]$ be the first Weyl algebra over a field F of characteristic zero. Now we consider the ring

$$R = \{(a_i)_{i=1}^{\infty} \mid a_i \in \text{Mat}_2(W) \text{ is eventually a constant upper triangular matrix}\},$$

where $\text{Mat}_2(W)$ denotes the full ring of 2×2 matrices over W . Then R is a semiprime ring of bounded index 2 whose prime ideals are maximal. But R is not π -regular. If every primitive factor ring of R is Artinian, then by Proposition 19, R is π -regular, which is a contradiction.

We conclude this paper with the following question.

Question 1. *If R is a right weakly π -regular ring of bounded index whose primitive factor rings are Artinian, then is R π -regular?*

Acknowledgements

The authors would like to thank Professor Jae Keol Park and Professor Y. Hirano for helpful comments and suggestions. The first named author was supported by the Academic Research Fund of Ministry of Education, Republic of Korea in 1997, Project No. BSRI-97-1432, while the second named author wishes to acknowledge the financial support of the Korea Research Foundation made in the Program Year 1997.

References

- [1] A. Badawi, On abelian π -regular rings, *Comm. Algebra* 25(4) (1997) 1009–1021.
- [2] G.F. Birkenmeier, H.E. Heatherly, E.K. Lee, Completely prime ideals and associated radicals, in: S.K. Jain, S.T. Rizvi (Eds.), *Proc. Biennial Ohio State-Denison Conf. 1992*, World Scientific, River Edge, NJ, 1993, pp. 102–129.
- [3] G.F. Birkenmeier, J.Y. Kim, J.K. Park, A connection between weak regularity and the simplicity of prime factor rings, *Proc. Amer. Math. Soc.* 122 (1994) 53–58.
- [4] G.F. Birkenmeier, J.Y. Kim, J.K. Park, Regularity conditions and the simplicity of prime factor rings, *J. Pure Appl. Algebra* 115 (1997) 213–230.
- [5] A.W. Chatters, C.R. Hajarnavis, *Rings with Chain Conditions*, Pitman Advanced Publishing Program, London, 1980.
- [6] F. Dischinger, Sur les anneaux fortment π -reguliers, *C.R. Acad. Sci. Paris Ser. A-B* 283 (1976) 571–573.
- [7] J.W. Fisher, R.L. Snider, On the von Neumann regularity of rings with regular prime factor rings, *Pacific J. Math.* 54(1) (1974) 135–144.
- [8] V. Gupta, Weakly π -regular rings and group rings, *Math. J. Okayama Univ.* 19 (1977) 123–127.
- [9] Y. Hirano, Some studies on strongly π -regular rings, *Math. J. Okayama Univ.* 20 (1978) 141–149.
- [10] Y. Hirano, Some characterizations of π -regular rings of bounded index, *Math. J. Okayama Univ.* 32 (1990) 97–101.
- [11] P. Menal, On π -regular rings whose primitive factor rings are Artinian, *J. Pure Appl. Algebra* 20 (1981) 71–78.
- [12] E.C. Posner, Prime rings satisfying a polynomial identity, *Proc. Amer. Math. Soc.* 11 (1960) 180–183.
- [13] H.H. Storrer, Epimorphismen von kommutativen Ringen, *Comment Math. Helv.* 43 (1968) 378–401.
- [14] S.H. Sun, Noncommutative rings in which every prime ideal is contained in a unique maximal ideal, *J. Pure Appl. Algebra* 76 (1991) 179–192.
- [15] Xue Yao, Weakly right duo rings, *Pure Appl. Math. Sci.* XXI(1-2) (1985) 19–24.
- [16] H.P. Yu, On quasi-duo rings, *Glasgow Math. J.* 37 (1995) 21–31.